

Electrophysiological Parameters in the Electrical Propagation During Atrial Fibrillation: A Population of Models Study

A Simon^{1,2}, A Liberos¹, I Hernandez-Romero¹, A Bueno-Orovio³, M Rodrigo⁴, MS Guillem⁴, F Atienza¹, F Fernandez-Aviles¹, B Rodriguez³, AM Climent¹

¹Cardiology Department, Hospital GU Gregorio Marañón, IiSGM, CIBERCV, Madrid, Spain

²Universidad Carlos III de Madrid, Spain

³Department of Computer Science, University of Oxford, Oxford, UK

⁴ITACA, Universitat Politècnica de Valencia, Valencia, Spain

Abstract

Cardiac tissue conduction velocity (CV) is a relevant variable in the maintenance and stability of atrial fibrillation (AF). The main goal of this work is to investigate modulation of CV in human chronic AF by inter-subject variability in ionic channel properties, cellular environmental factors and tissue coupling, by using a population of mathematical models. A monodomain tissue model was developed on the basis of the Skibsbjerg human atrial cell model. In this model, nine parameters are varied leading to a total number of 500 tissue models (8x156 cells). Among these simulations, 126 models are found within the experimental physiological range of biomarkers obtained from 149 chronic AF patients. The resulting population of 126 tissue models covers the variability in the biomarkers observed in the experimental recordings. The specific balance between sodium current and diffusion coefficient modulates the CV, critical for arrhythmia inducibility. The developed population of tissue models establishes the basis for further research in ionic mechanisms that facilitate AF maintenance and in-silico evaluation of personalised drug treatments.

1. Introduction

Atrial fibrillation (AF) is the most prevalent cardiac arrhythmia encountered in the clinical practice [1]. Moreover, the efficacy of current therapies to restore sinus rhythm is limited, especially in patients suffering chronic AF [1]. A deep understanding of the mechanisms responsible for the initiation and maintenance of the arrhythmia is essential for the increase of treatment efficacy [2].

Modifications in electrophysiological characteristics, known as atrial remodeling, facilitate reentrant stability thanks to the shortening of action potential (AP) and the reduction in conduction velocity (CV). According to other biological properties, this remodeling process is highly dependent on the genetics and clinical scenario of each patient, which explains the differences in the

efficacy of treatments depending on the patient [2, 3].

Populations of models aim to overcome and prevent from unexpected responses and secondary effects associated with electrophysiological inter-subject variability during the implementation of new drugs [2]. Previous inter-subject variability studies have focused on the differences observed in representative biomarkers at the cellular level but, when studying mechanisms responsible for the initiation and maintenance of AF, it is crucial to consider the effect of neighboring cells and alterations in CV. The organization and structure of atrial tissue has been associated with atrial conduction abnormalities, presenting a substrate for atrial reentry [4].

In this work, we propose a population of 2D atrial tissue models, calibrated against human recordings in chronic AF (cAF), to investigate modulation of human atrial CV by inter-subject variability in ionic current densities, environmental factors, and tissue coupling.

2. Materials and methods

2.1. Experimental dataset and biomarkers

A set of experimental recordings were used to calibrate the population of models. Specifically, measurements on right atrial appendages from 149 patients diagnosed with cAF were employed [2,5]. A total of six biomarkers were used to quantify variability in APs (Figure 1B): AP duration at 20, 50, and 90% of repolarization (APD20, APD50, APD90 respectively), AP amplitude (APA), resting membrane potential (RMP) and AP plateau potential at 20% of APD90 (V20). The maximum and minimum values of these biomarkers at a pacing frequency of 1Hz are presented in Table 1.

2.2. Electrophysiological cellular model

The Skibsbjerg 2016 model was implemented and used in this work [6], whose schematic is depicted in Figure 1A. The model provides a detailed description of the transmembrane ionic currents and intracellular calcium handling, as an extension of the Koivumaki model [7].

[Escriba aquí]

stimuli. A cardiac simulation GP-GPU platform was used to perform the mathematical simulations. Biomarkers for the 500 tissue models were evaluated, and only those satisfying all the experimental constraints were included in the population of models.

Partial correlation was used to find correlations between electrophysiological parameters and biomarkers, after accounting for the linear effects of one or more additional variables [2, 3].

3. Results

3.1 Generation of the population of models

126 of the total 500 models (25.2%) presented all the biomarkers within the experimental range (Table 1). Only this set of experimentally calibrated population can be considered physiologically acceptable. The physiological population of models and experimental data are presented in Figure 2. Experimental recordings (orange dots) are compared with model predictions (blue dots) showing a good agreement between them. In addition, baseline AF model predictions are shown with green dots.




Figure 2. Population of tissue models calibration and distribution of CVs. Panels A, B and C show the values of the biomarkers $-V_{20}$, APD_{20} , APD_{50} , APD_{90} , RMP and APA for the experimentally calibrated models. Blue marks correspond to the ‘physiological population’ ($N=126$). Orange marks correspond with experimental recordings. Baseline values are marked in green. Upper and lower bounds for 1 Hz biomarkers are marked in light blue. Panel D illustrates the distribution in CV of the physiological acceptable models.

[Escriba aquí]

Figure 2D shows the distribution of CV values in the physiological population. The obtained CVs ranges from 18.06 cm/s to 100.77 cm/s, covering similar values than those presented in previous studies [9].

The results illustrate how the population of models in tissue is able to reproduce the inter-subject variability found in patients (every experimental value has a neighboring model). Furthermore, the biomarkers corresponding to the baseline AF model were located inside experimental constraints.

Box plots in Figure 3A show the range of the varied parameters and the median physiological values. Edges of the boxes represent the 1st and 3rd quartiles. Note that most of the median values do not coincide with the baseline model value (g_{Na} , I_{NaK} , g_{CaL} , I_{KCa} , $[Na^+]_o$ and D) that is represented by the green line. These results indicate that the original model is not representative of median values. A wide range of values for the diffusion coefficient, ionic conductances and concentrations is covered. Most of the parameters admitted to be varied from -50% up to +90%, or even 100% of the basal value.




Figure 3. Box plot and partial correlation analysis. Box plot of Panel A shows the range and mean value of the varied parameters. Panel B shows partial correlation results where each correlation represents the effect of a specific parameter on each biomarker. Positive and negative correlations are marked in red and blue, respectively.

The results obtained after partial correlation analysis are presented in Figure 3B. Each correlation represents the effect of a specific parameter on each biomarker and the intensity of the color shows the intensity of the

correlation. Positive and negative correlations are marked in red and blue, respectively. Most of the parameters have a strong effect on various biomarkers. However, I_{KCa} and $[Na^+]_o$ present weaker correlations with the biomarkers. Regarding CV, the presented study demonstrates a strong direct correlation of g_{Na} and D with CV, whereas weaker correlations are established with I_{NaK} , g_{kl} and $[Na^+]_o$.

On the other hand, D slightly affects V20 and RMP but no correlation was found with the rest of biomarkers. The analysis also concludes that an increase in sodium conductivity is directly related to APA and CV, and inversely related with APD20, APD50 and APD90. APD90 was found to be inversely related to potassium and sodium currents while it is directly related with g_{CaL} . Finally, $[K^+]_o$ was found to be strongly related with RMP and inversely related to APA and APD90.

4. Discussion

In this study, the effects of tissue inter cellular coupling on atrial propagation have been investigated in a subset of parameters, based in previous populations of models analysis [2, 3, 5] with other parameters underlying electrical propagation. The results obtained in this work are consistent with previous studies carried out at cellular level. Taking this as a sign of fidelity, the tissue level model has been used to analyze the problem in a deeper way by taking into account other features like CV and electrical propagation.

As a result, we have demonstrated that populations of models are affected by inter-cellular coupling that, together with the sodium current, modulates CV. Consequently, alteration of these parameters is highly relevant when studying the maintenance and stability of AF. Specifically, CV is predominantly governed by the specific balance between D and g_{Na} . However, although g_{kl} is strongly correlated with RMP, we did not observe a strong relation between g_{kl} and CV at the examined pacing frequency. Furthermore, the included parameter I_{KCa} , does not strongly affect any biomarker. Finally, it will be important for future studies to consider the population at different frequencies to ensure that the population reproduces rate dependent properties.

5. Conclusion

The use of populations of models at the cardiac tissue level can help in future investigations in the identification of the different mechanisms underlying AF initiation and termination, and thus a better classification of patients according to their most prominent mechanism. The development of populations of models accounting for diffusion properties can serve as a basis for further

[Escriba aquí]

research in the evaluation of personalized anti-AF pharmacological therapies.

Acknowledgements

This work has been possible thanks to the financing of the Instituto de Salud Carlos III and Ministerio de Ciencia e Innovación.

References

- [1] Guillem MS, Climent AM, Rodrigo M, Fernández-Aviles F, Atienza F, Berenfeld O. Presence and stability of rotors in atrial fibrillation: Evidence and therapeutic implications. *Cardiovasc Res* 2016;109(4):480–92.
- [2] Liberos A, Bueno-Orovio A, Rodrigo M, Ravens U, Hernandez-Romero I, Fernandez-Aviles F, et al. Balance between sodium and calcium currents underlying chronic atrial fibrillation termination: An in silico inter-subject variability study. *Heart Rhythm* 2016;13(12):2358-65.
- [3] Britton OJ, Bueno-Orovio A, Van Ammel K, Lu HR, Towart R, Gallacher DJ, et al. Experimentally calibrated population of models predicts and explains intersubject variability in cardiac cellular electrophysiology. *Proc Natl Acad Sci USA* 2013;110(23):E2098-105.
- [4] Nattel S, Maguy A, Bouter SLE, Yeh Y. Arrhythmogenic ion-channel remodeling in the heart: Heart failure, myocardial infarction, and atrial fibrillation. *Physiol Rev* 2007;87:425–56. doi:10.1152/physrev.00014.2006.
- [5] Sánchez C, Bueno-Orovio A, Wettwer E, Loose S, Simon J, Ravens U, et al. Inter-subject variability in human atrial action potential in sinus rhythm versus chronic atrial fibrillation. *PLoS One* 2014; 9:e105897.
- [6] Skibsbjerg L, Jespersen T, Christ T, Maleckar MM, Brink J Van Den, Tavi P, et al. Refractoriness in human atria: Time and voltage dependence of sodium channel availability. *J Mol Cell Cardiol* 2016;101:26–34.
- [7] Koivumäki JT, Seemann G, Maleckar MM, Tavi P. In silico screening of the key cellular remodeling targets in chronic atrial fibrillation. *PLoS Comput Biol* 2014;10(5):e1003620.
- [8] Clayton RH, Panfilov AV. A guide to modelling cardiac electrical activity in anatomically detailed ventricles. *Prog Biophys Mol Biol* 2008;96(1–3):19–43.
- [9] Wilhelms M, Hettmann H, Maleckar MM, Koivumäki JT, Dössel O, Seemann G. Benchmarking electrophysiological models of human atrial myocytes. *Front Physiol* 2013;3:487.

Address for correspondence.

Andreu M. Climent

Investigación Sanitaria Gregorio Marañón Hospital GU Gregorio Marañón. Edificio Materno Infantil.

c/ O'Donnell 48, 28009, Madrid, Spain

andreu.climent@gmail.com