

# Effects of CaMKII Regulation on Atrial Action Potential Under Oxidative Stress Condition

Haibo Sui<sup>1</sup>, Qince Li<sup>1\*</sup>, Kuanquan Wang<sup>1</sup>, Funebi Francis Ijebu<sup>1</sup>, Yongfeng Yuan<sup>1</sup>, Henggui Zhang<sup>1,2,3,4</sup>

<sup>1</sup> School of Computer Science and Technology, Harbin Institute of Technology, Harbin, Heilongjiang, China

<sup>2</sup> School of Physics and Astronomy, The University of Manchester, Manchester, UK

<sup>3</sup> Space Institute of Southern China (Shenzhen), Shenzhen, China

<sup>4</sup> Key Laboratory of Medical Electrophysiology, Ministry of Education, Collaborative Innovation Center for Prevention and Treatment of Cardiovascular Disease/Institute of Cardiovascular Research, Southwest Medical University, Luzhou, China

## Abstract

*Previous studies have demonstrated that oxidative stress is closely associated with cardiac arrhythmias via altering electrical activity and intracellular calcium dynamics of cardiac myocytes. The present study developed a human atrial cell model including effects of oxidative stress by incorporating reactive oxygen species (ROS)-induced CaMKII activation and its downstream effects on different ionic channels. The CaMKII dynamics was mimicked by a novel 6-state Markov chain model, including both autophosphorylation and oxidation pathways of CaMKII activation. Simulation results show that highly activated CaMKII by massive ROS phosphorylated L-type  $Ca^{2+}$  current ( $I_{CaL}$ ) and elevated intracellular calcium concentration, which eventually resulted in calcium overload in sarcoplasmic reticulum in the condition of oxidative stress. Meanwhile, calcium overload resulted in increase of calcium release and cytoplasmic calcium concentration, which triggered afterdepolarizations (DADs) by increasing the calcium extrusion via the  $Na^+$ - $Ca^{2+}$  exchanger (NCX) current. This study revealed effects of massive ROS on calcium cycling in atrial myocytes and shed lights on mechanisms of cardiac arrhythmias induced by oxidative stress.*

## 1. Introduction

Several experiments have revealed that Calcium-calmodulin dependent kinase II (CaMKII) is a critical player in promotion of cardiovascular disease and arrhythmia phenotypes [1]. CaMKII can regulate membrane ion channels and  $Ca^{2+}$ -handling proteins in cardiac myocytes and is the central mediator of excitation-

contraction coupling. CaMKII is activated by binding of  $Ca^{2+}$ -bound calmodulin and can keep persistent activity through autophosphorylation and oxidation. CaMKII activation under oxidation has become a biomarker and proarrhythmic signal for connecting ROS and arrhythmia.

Experiment has shown that increased CaMKII expression results in hyperphosphorylation of RyR [2] that can promote diastolic sarcoplasmic reticulum  $Ca^{2+}$  release and leak with the potential to trigger after-depolarization (DAD). However, the effect of CaMKII on atrial action potential under oxidative stress condition is not completely understood. In this study, we developed a computational model of human atrial cell including CaMKII oxidative activation to investigate influences of CaMKII on human atrial electrophysiology.

## 2. Methods

CaMKII activation model developed by Zhang et al [3] was incorporated into a human atrial model developed by Grandi et al [4] to simulate CaMKII activation by autophosphorylation and oxidation. The parameters of the CaMKII autophosphorylation pathway were based on the model developed by Chiba et al [5]. The parameters of oxidation pathway were fitted based on the experimental data measured by Erickson et al [6]. To simulate the effects of CaMKII on ion channels, we modified the equations of corresponding membrane currents using the method proposed by O'hara et al [7]. In this approach, the effect of CaMKII on a single current was simulated using Hill equation and, therefore, the amplitude and/or time constant of ion current were modified based on experiments.

RyR and PLB are very important regulatory protein on the SR to control calcium cycling. RyR can be phosphorylated by activated CaMKII, increasing RyR

opening probability and SR leak. PLB phosphorylated by activated CaMKII can reduce SERCA inhibition, increasing SR pump calcium sensitivity during diastole. Thus, we incorporated RyR and PLB model developed Soltis et al [8]. The rate constants which controlling RyR opening probability and SR leak was modified under CaMKII activation. The altered rate constant increased calcium sensitivity of RyR and amplified RyR opening. Also, the equation of SR leak, which is associated with RyR opening at the diastolic phase, was modified under CaMKII activation. The Thr17 of PLB can be phosphorylated by CaMKII to increase SERCA pump calcium sensitivity which means decreasing the forward pump rate. This phenomenon was mimicked by modifying half maximal saturation constant of SERCA.

To investigate the effect oxidative CaMKII activation on the electrophysiology of cells, a dynamic stimulation process was used in the simulation. The stimulation frequency was selected as 0.5Hz, 1Hz and 2Hz to investigate the frequency dependence of CaMKII activation. Action potential were stimulated by a stimulus current of  $-12.5\text{pA/pF}$  for 0.5ms. The time step of simulation was 0.1ms. Under normal conditions, the level of  $\text{H}_2\text{O}_2$  in human blood is about  $\sim 35\mu\text{M}$ . When ischemic reperfusion injury or heart failure occurred, the level of ROS could be increased by 100 times. Therefore, the oxidative stress was simulated using  $200\mu\text{M}$   $\text{H}_2\text{O}_2$  in the model.

### 3. Results

#### 3.1. Frequency dependent CaMKII activation

Previous experiment has shown that the activation rate and maximum activation level of CaMKII increase as frequency increases [9]. The frequency-dependent characteristics of CaMKII were investigated in the model without ROS. Fig. 1 shows the simulation results of activation level of CaMKII at 0.5Hz, 1Hz and 2Hz (Fig. 1A). Meanwhile, the activation level of RyR2 in the same condition (Fig. 1B) was recorded as well. The results showed that the slope of CaMKII activation curves increased with enlarging frequency. The RyR activation level showed a similar behaviour with increasing frequency, indicating a positive correlation between CaMKII activation and RyR opening probability. In addition, the maximum percentage of CaMKII activation and RyR phosphorylation increased with increasing pacing rate (Fig. 1 A and B). These results agreed with previous RyR phosphorylation model developed [10], demonstrating that CaMKII activation has a significantly influence on SR  $\text{Ca}^{2+}$  release and leak.




Figure 1. Simulation results of CaMKII activation and RyR phosphorylation from initial condition to stable state at 0.5Hz, 1Hz and 2Hz in the 60s.

#### 3.2. $\text{H}_2\text{O}_2$ induced DADs

According to simulation results, a DAD was triggered in the action potential in conditions of elevated oxidative stress (Fig. 2). Due to the frequency-dependent CaMKII activation,  $\text{H}_2\text{O}_2$ -induced DADs showed the rate dependence as well, which indicated more significant DADs were induced at higher frequency of stimulation. In this study, we chose to intercept part of the periodic data in the case of 0.5Hz to analysis the generation of DAD.

We inhibited NCX current at 0.5Hz with the presence  $\text{H}_2\text{O}_2$  and to examine its contribution to  $\text{H}_2\text{O}_2$  induced DAD (Fig. 3). The result shows that DAD did not occur with inhibiting NCX current, which is consistent with the experimental findings [11]. Since the mode of NCX current is determined by the concentration gradient on both sides of membrane, the increase of intracellular calcium promoted the forward transport of NCX. Meanwhile, CaMKII did not directly change NCX. Therefore, the release of calcium from SR to the cytoplasm (Fig. 2F-H) during phase 4 of AP induced the inflow of NCX, which leads to depolarization of the membrane potential and production of DAD.




Figure 2. Simulated AP (A), CaMKII activation fraction (B) and CaMKII regulated current (C, D, E) and  $Ca^{2+}$  flux (F, G, H) under both control (Black curves) and oxidative stress condition (red curves).

CaMKII can regulate numerous ion channels and proteins in the membrane and intracellular, including  $I_{CaL}$ ,  $I_{Na}$  and  $Ca^{2+}$  cycling related proteins. The simulation results showed that oxidative stress increased CaMKII activation, increased  $I_{CaL}$  and altered  $Ca^{2+}$  cycling (Fig. 2). Therefore, we simulated the action potential under inhibition of CaMKII effect on single ion channel. In Fig. 4A, DAD occurrence has been greatly suppressed when inhibiting CaMKII effect on  $I_{CaL}$ . In addition, the DAD amplitude is reduced after inhibiting CaMKII effect on SR calcium release or pump (Fig. 4B and C). But inhibiting CaMKII effect on SR calcium leak makes the occurrence of repetitive DADs (Fig. 4D). This case indirectly indicated that SR  $Ca^{2+}$  overload causes the generation of DADs. CaMKII activation has been shown to increase RyR opening probability and SR leakage. Inhibiting CaMKII effect on SR leak reduced the amount of leaked calcium which means it is easier for accumulate of calcium concentration in SR to overload, and leading to repetitive DAD phenomenon. CaMKII indirectly regulates SR calcium release, pump and leak in the  $Ca^{2+}$  cycling by phosphorylating of RyR. It was shown that DAD was eliminated when the level of RyR phosphorylation was clamped at normal level (Fig. 4E). The simulation results showed that DAD was suppressed obviously. From the above results, it indicated that CaMKII regulated  $I_{CaL}$  and RyR could be important targets for eliminating the abnormal calcium cycling and DAD.




Figure 3. In oxidative stress condition, simulated AP in the 0.5Hz with NCX block. Note that no DAD occur in the NCX block condition.




Figure 4. Simulated AP and SR  $Ca^{2+}$  concentration with inhibition of CaMKII effect on  $I_{CaL}$  (A, F), SR uptake (B, G), SR release (C, H), and SR leak (D, I) as well as with clamped RyR phosphorylation (E, J) in oxidative stress condition.




Figure 5. In oxidative stress condition, simulated AP and SR  $Ca^{2+}$  concentration under the conditions of CaMKII affecting  $I_{CaL}$  (A), RyR (B) only and both (C) respectively.

In Fig. 5, with CaMKII regulating  $I_{CaL}$  and RyR alone, there was no obvious DAD in action potential. However, when CaMKII regulated both  $I_{CaL}$  and RyR, there was an obvious DAD abnormality in action potential.  $I_{CaL}$  was not able to sufficiently increase the concentration of calcium in cytoplasm to induce DAD. The combined effect of  $I_{CaL}$  and RyR contributed to the elevated calcium concentration in cytoplasm and caused DAD generation.

In conclusion, the oxidative pathway of CaMKII significantly increased CaMKII activation and the RyR opening probability in oxidative stress condition. PLB phosphorylated by activated CaMKII reduced the inhibitory effect on SR pump, which in turn increased calcium uptake capacity of SR pump. These effects increased the calcium concentration of SR and led to SR calcium overload finally. Calcium overload produced a spontaneous calcium in phase 4, stimulating the inflow of NCX current on the cell membrane and leading to the occurrence of DAD.

#### 4. Conclusion

In this study, we developed a human atrial computational model including CaMKII oxidation and revealed the mechanism of DAD induced under oxidative stress condition. The simulation results indicated that  $I_{CaL}$  and RyR regulated by CaMKII in oxidative stress condition primarily contributed to DAD generation, as either of them regulated by CaMKII is insufficient to cause  $H_2O_2$ -induced DAD.

#### Acknowledgements

The work is supported by the National Nature Science Foundation of China (NSFC) (No. 61601143, 81770328,

61572152, and 61571165), Heilongjiang and China Postdoctoral Science Foundation under Grant nos.2015M581448 (to QL), the Natural Science Foundation of Shandong Province (No. ZR2017MF051), the Science Technology and Innovation Commission of Shenzhen Municipality (No. JSGG20160229125049615 and JCYJ20151029173639477).

#### References

- [1] Mesubi O O, Anderson M E. Atrial remodelling in atrial fibrillation: CaMKII as a nodal proarrhythmic signal. *Cardiovascular Research*, 2016, 109(4): 542-557.
- [2] Dobrev D, Wehrens X H. Calcium-mediated cellular triggered activity in atrial fibrillation. *Journal of Physiology*, 2017, 595(12).
- [3] Zhang S, Li Q, Zhou L, et al. Development of a novel Markov chain model for oxidative-dependent CaMKII activation. *Computing in Cardiology Conference. IEEE*, 2015:881-884.
- [4] Grandi E, Pandit S V, Voigt N, et al. Human atrial action potential and  $Ca^{2+}$  model: sinus rhythm and chronic atrial fibrillation. *Circulation Research*, 2011: CIRCRESAHA. 111.253955.
- [5] Chiba H, Schneider N S, Matsuoka S, et al. A Simulation Study on the Activation of Cardiac CaMKII  $\delta$ -Isoform and Its Regulation by Phosphatases. *Biophysical Journal*, 2008, 95(5):2139-2149.
- [6] Erickson J R, Joiner M L, Guan X, et al. A dynamic pathway for calcium-independent activation of CaMKII by methionine oxidation. *Cell*, 2008, 133(3):462-474.
- [7] O'Hara T, Virá g L, Varrá s A, et al. Simulation of the undiseased human cardiac ventricular action potential: model formulation and experimental validation. *Plos Computational Biology*, 2011, 7(5):e1002061.
- [8] Soltis A R, Saucerman J J. Synergy between CaMKII substrates and  $\beta$ -adrenergic signaling in regulation of cardiac myocyte  $Ca^{2+}$  handling. *Biophysical Journal*, 2010, 99(7):2038-2047.
- [9] Dupont G, Houart G, De K P. Sensitivity of CaM kinase II to the frequency of  $Ca^{2+}$  oscillations: a simple model. *Cell Calcium*, 2003, 34(6):485-497.
- [10] Saucerman J J, Healy S N, Belik M E, et al. Proarrhythmic consequences of a KCNQ1 AKAP-binding domain mutation: computational models of whole cells and heterogeneous tissue. *Circulation Research*, 2004, 95(12):1216.
- [11] Xu L, Li X Y, Ren S, et al. Analysis of Na/Ca exchanger (NCX) function and current in murine cardiac myocytes during heart failure. *Molecular Biology Reports*, 2012, 39(4):3847-3852.

Address for correspondence

Qince Li  
Xinjishu Building 901,  
Harbin Institute of Technology  
Xidazhi Street, Nangang District,  
Harbin,  
China

qinceli@hit.edu.cn